

Reported Cases of Syphilis, New Mexico, 1970-2007

New Mexico Cases

Primary & Secondary Syphilis

Year	U.S. Cases	U.S. Rate* per 100,000	New Mexico Cases	New Mexico Rate* per 100,000	New Mexico Ranking for P&S Cases	Early Latent Syphilis Cases	Congenital Syphilis Cases	Late Syphilis Cases	All Stages Syphilis
1970	21,982	10.9	125	12.3		119	no data	420	664
1971	23,783	11.6	162	15.5		132	8	236	538
1972	24,429	11.8	109	10.2		147	16	203	475
1973	24,825	11.8	88	8.0		97	11	344	540
1974	25,385	12.0	97	8.6		86	4	328	515
1975	25,561	12.0	154	13.4		134	10	223	521
1976	23,731	11.0	147	12.6		129	7	207	490
1977	20,399	9.4	102	8.6		78	4	182	366
1978	21,656	9.8	89	7.3		55	7	126	277
1979	24,874	11.2	95	7.8		64	3	71	233
1980	27,204	12.1	115	8.9		105	0	52	272
1981	31,266	13.7	132	10.2		116	4	100	352
1982	33,613	14.6	203	15.6		118	1	62	384
1983	32,698	14.1	185	13.5		163	1	80	429
1984	28,607	12.1	99	7.0	20th	94	0	68	261
1985	27,131	11.4	127	8.8	16th	103	0	88	318
1986	27,667	11.5	78	5.3	21st	86	0	112	276
1987	35,585	14.7	59	4.0	25th	61	1	208	329
1988	40,474	16.7	75	5.0	27th	98	0	290	463
1989	45,826	18.6	31	2.1	32nd	50	0	109	190
1990	50,578	20.3	51	3.4	31st	60	4	69	184
1991	42,950	17.0	41	2.6	37th	44	5	78	168
1992	34,009	13.3	44	2.8	34th	36	3	48	131
1993	26,527	10.2	32	2.0	35th	59	2	87	180
1994	20,641	7.8	15	0.9	36th	22	2	97	136
1995	16,543	6.2	13	0.8	36th	30	1	88	132
1996	11,405	4.2	7	0.4	40th	10	1	53	71
1997	8,556	3.1	14	0.8	30th	21	1	82	118
1998	7,007	2.5	24	1.3	23rd	25	3	51	103
1999	6,617	2.4	15	0.8	24th	11	2	67	95
2000	5,979	2.1	16	0.9	23rd	30	3	52	101
2001	6,103	2.1	20	1.1	24th	25	5	34	84
2002	6,862	2.4	39	2.0	18th	30	7	44	120
2003	7,177	2.5	73	3.8	6th	61	6	76	216
2004	7,980	2.7	82	4.2	5th	80	3	97	262
2005	8,724	3.0	56	2.8	14th	56	6	70	188
2006	9,642	3.3	78	3.9	10th	91	7	73	249
2007**	11,301	3.8	46	2.2	23rd	65	6	61	178

Data as of 7/15/2008

* Rates are estimates based on data from the U.S. Census Bureau and the University of New Mexico, Bureau of Business and Economic Research (www.unm.edu/~bber)

** For 2007, all U.S. numbers and rates as well as the New Mexico ranking are preliminary

During 2007 and 2008, the New Mexico STD Program conducted extensive quality assurance review of STD case investigations from 1999 through 2007. These QA activities have resulted in identifying hundreds of STD cases (mostly chlamydia and most from 2004-2006) that had not been entered into the STD database, hundreds of miscodes for Race/Ethnicity (mostly cases among Native American incorrectly coded as Other) and numerous incorrect ages and county of residence. These errors have been fixed. (See note below about errors that could not be corrected) As result, the numbers of cases on these tables will differ from what had been posted on this website prior to September 2008.