



HIV Infection and AIDS among Adults and Adolescents, New Mexico, 2012

	Newly Diagnosed in 2012*			Persons Living with HIV or AIDS in New Mexico								
	N	%	Rate [†]	HIV			AIDS			Total		
				N	%	Rate [†]	N	%	Rate [†]	N	%	Rate [†]
Total	119	100	7.0	1132	100	66.2	1768	100	103.4	2900	100	169.5
Sex at Birth												
Male	108	91%	12.9	964	85%	114.9	1,573	89%	187.5	2,537	87%	302.4
Female	11	9%	1.3	168	15%	19.3	195	11%	22.4	363	13%	41.7
Race/Ethnicity												
American Indian/Alaska Native	21	18%	14.7	101	9%	70.9	131	7%	92.0	232	8%	162.9
Asian/Pacific Islander	0	0%	0	6	1%	22.9	10	1%	38.2	16	1%	61.1
Black	2	2%	5.9	89	8%	261.4	97	5%	284.9	186	6%	546.3
Hispanic	66	55%	8.8	515	45%	68.7	774	44%	103.2	1289	44%	171.9
White	28	24%	3.7	407	36%	53.7	723	41%	95.4	1130	39%	149.1
Multi-race	2	2%	CNC	14	1%	CNC	33	2%	CNC	47	2%	CNC
Age[‡]												
13-24	20	17%	5.7	56	5%	16.1	16	1%	4.6	72	2%	20.7
25-34	41	34%	15.2	245	22%	91.0	132	7%	49.0	377	13%	140.0
35-44	28	24%	11.2	294	26%	118.1	363	21%	145.8	657	23%	263.8
45-54	25	21%	8.7	330	29%	114.9	713	40%	248.3	1043	36%	363.3
55+	5	4%	0.9	207	18%	37.2	544	31%	97.7	751	26%	134.9
Region[§]												
Northwest	18	15%	9.9	107	9%	59.0	148	8%	81.6	255	9%	140.5
Northeast	22	18%	8.8	197	17%	78.6	358	20%	142.8	555	19%	221.4
Metro	50	42%	6.7	541	48%	72.9	919	52%	123.8	1,460	50%	196.7
Southeast	5	4%	2.2	54	5%	23.4	104	6%	45.1	158	5%	68.5
Southwest	24	20%	7.9	233	21%	76.4	239	14%	78.3	472	16%	154.7
Transmission Category												
Men who have sex with men (MSM)	71	60%	CNC	664	59%	CNC	1095	62%	CNC	1759	61%	CNC
Intravenous Drug Use (IDU)	8	7%	CNC	88	8%	CNC	161	9%	CNC	249	9%	CNC
MSM/IDU	10	8%	CNC	72	6%	CNC	184	10%	CNC	256	9%	CNC
High-Risk-Hetrosexual	16	13%	CNC	137	12%	CNC	181	10%	CNC	318	11%	CNC
Other	0	0%	CNC	2	0%	CNC	12	1%	CNC	14	0%	CNC
Perinatal Exposure	0	0%	CNC	4	0%	CNC	8	0%	CNC	12	0%	CNC
No Risk Identified	14	12%	CNC	165	15%	CNC	127	7%	CNC	292	10%	CNC

Footnotes

*Persons age 13 or older who were newly diagnosed while residing in New Mexico during 2012
[†]Rates per 100,000 based on Bureau of Business and Economic Research Geospatial and Population Studies Program population estimates for 2011.
[‡]For new diagnoses, age at time of diagnosis. For persons living with HIV or AIDS, current age.
[§]For new diagnoses, residence at time of diagnosis. For persons living with HIV or AIDS, current residence.
 CNC means Can Not Calculate - population data are unavailable.

Cautions

N < 20: Rate calculations for small numbers may be statistically unreliable. Fluctuations in the associated rates may be due in part to statistical unreliability rather than true change.
 Region: New Mexico Department of Health constructed new health regions in 2012. Regional comparison to previous years is not recommended. A map depicting New Mexico's health regions is available for download from the New Mexico Department of Health at <http://ibis.health.state.nm.us/image/HealthRegions2012.png>.