

 HIV Surveillance Program Epidemiology and Response Division New Mexico Department of Health	HIV Infection and AIDS among Adults and Adolescents, New Mexico 2016											
	Newly Diagnosed*			Persons Living with HIV or AIDS†								
				HIV			AIDS			Total		
	N	%	Rate‡	N	%	Rate‡	N	%	Rate‡	N	%	Rate‡
Total	121	100.0	6.9	1,507	100.0	86.2	1,942	100.0	111.1	3,449	100.0	197.3
Sex at birth												
Male	110	90.9	12.8	1,305	86.6	151.4	1,734	89.3	201.2	3,039	88.1	352.6
Female	11	9.1	1.2	202	13.4	22.8	208	10.7	23.5	410	11.9	46.3
Race/Ethnicity												
American Indian/Alaska Native	20	16.5	13.9	136	9.0	94.3	151	7.8	104.7	287	8.3	199.0
Asian/Pacific Islander	0	0.0	0.0	11	0.7	35.9	7	0.4	22.9	18	0.5	58.8
Black	4	3.3	10.6	112	7.4	297.9	99	5.1	263.3	211	6.1	561.2
Hispanic	71	58.7	8.8	739	49.0	91.8	872	44.9	108.3	1,611	46.7	200.1
White	25	20.7	3.4	489	32.5	66.9	776	40.0	106.2	1,265	36.7	173.1
Multi-race	1	0.8	CNC	20	1.3	CNC	37	1.9	CNC	57	1.7	CNC
Age, years§												
13–24	27	22.3	7.9	81	5.4	23.6	14	0.7	4.1	95	2.8	27.7
25–34	48	39.7	17.0	351	23.3	124.0	121	6.2	42.7	472	13.7	166.7
35–44	25	20.7	10.2	341	22.6	139.2	312	16.1	127.4	653	18.9	266.6
45–54	13	10.7	5.1	376	25.0	148.0	680	35.0	267.7	1,056	30.6	415.7
55+	8	6.6	1.3	358	23.8	57.5	815	42.0	130.9	1,173	34.0	188.5
Region¶												
Northwest	17	13.2	9.5	160	10.6	89.4	175	9.0	97.7	335	9.7	187.1
Northeast	11	8.3	4.3	249	16.5	97.6	380	19.6	148.9	629	18.2	246.5
Metro	68	47.1	8.9	752	49.9	98.0	995	51.2	129.7	1,747	50.7	227.7
Southeast	8	5.8	3.4	93	6.2	39.2	113	5.8	47.6	206	6.0	86.8
Southwest	17	13.2	5.5	253	16.8	81.8	279	14.4	90.3	532	15.4	172.1
Transmission Category												
Men who have sex with men (MSM)	65	53.7	CNC	919	61.0	CNC	1,220	62.8	CNC	2,139	62.0	CNC
Intravenous Drug Use (IDU)	7	5.8	CNC	103	6.8	CNC	176	9.1	CNC	279	8.1	CNC
MSM/IDU	9	7.4	CNC	113	7.5	CNC	180	9.3	CNC	293	8.5	CNC
High-Risk-Heterosexual	6	5.0	CNC	164	10.9	CNC	186	9.6	CNC	350	10.15	CNC
Other	0	0.0	CNC	2	0.1	CNC	11	0.6	CNC	13	0.38	CNC
Perinatal Exposure	0	0.0	CNC	4	0.3	CNC	8	0.4	CNC	12	0.4	CNC
No Risk Identified	12	9.9	CNC	122	8.1	CNC	107	5.5	CNC	229	6.6	CNC
No Risk Reported	22	18.2	CNC	80	5.3	CNC	54	2.8	CNC	134	3.9	CNC

Footnotes

NOTE: Distribution of prevalent HIV and AIDS cases amongst the total Persons Living with HIV or AIDS in New Mexico is 43.7% and 56.3 %, respectively.
 NOTE: CNC means Can Not Calculate - population data is unavailable.
 NOTE: Percentages across characteristics within a section of this table may not add to 100.0% as a result of rounding.
 *All people who were newly diagnosed with Stage 1 through 3 HIV infection while residing in New Mexico during 2016; age 13 & older.
 † All people newly or previously diagnosed with Stage 1 through 2 or Stage 3 HIV infection and residing in New Mexico during 2016; age 13 & older.
 ‡ Rates per 100,000 based on Bureau of Business and Economic Research Geospatial and Population Studies Program population estimates for 2016.
 § For new diagnoses, age at time of diagnosis. For persons living with HIV or AIDS, current age.
 ¶ For new diagnoses, residence at time of diagnosis. For persons living with HIV or AIDS, current residence.

Cautions

N < 20: Rate calculations for small numbers may be statistically unreliable. Fluctuations in the associated rates may be due in part to statistical unreliability rather than true change.
 Region: New Mexico Department of Health constructed new health regions in 2012. Regional comparison to previous years is not recommended. A map depicting New Mexico's Health Regions is available for download from the New Mexico Department of Health at <https://ibis.health.state.nm.us/view/docs/HealthRegions2012.pdf>.
 Red "***" - Indicates ≥ 20% positive change from previous year, for values > 20.
 Green "++" - Indicates ≥ 20% negative change from previous year, for values > 20.